



Sleep

Some people may experience sleepiness or insomnia because of their antidepressant.




Weight Change

Some people may experience weight change. It is most likely to occur over six to twelve months and depends on your actual weight. The chart below is based on a 150 lb person.




Stopping Approach

Quitting your medicine all at once can make you feel sick, as if you had the flu (e.g. headache, dizziness, light-headedness, nausea or anxiety).



Sexual Issues

Some people may experience loss of sexual desire (libido) or loss of ability to reach orgasm because of their antidepressant.



Cost

Keep in Mind

Making Wiser Choices About Medicines

These figures are estimates and are for comparative reference only. Actual out-of-pocket costs vary over time, by pharmacy, insurance plan coverage, preparation and dosage.

Less ← → More

Category	Medicine (Brand Name)	Cost / month
SSRIs	Citalopram (Celexa®)	\$4 / month – Super-stores drug program
	Escitalopram (Lexapro®)	\$110 / month
	Fluoxetine (Prozac®)	\$4 / month – Super-stores drug program
	Fluvoxamine (Luvox®)	\$80 / month
	Paroxetine (Paxil®)	\$4 / month – Super-stores drug program
	Sertraline (Zoloft®)	\$29 / month
SNRIs	Desvenlafaxine (Pristiq®)	\$147 / month – No generic available
	Duloxetine (Cymbalta®)	\$154 / month – No generic available
	Venlafaxine (Effexor®)	\$130 / month
Others	Bupropion (Wellbutrin®)	\$100 / month
	Mirtazapine (Remeron®)	\$50 / month
TCAs	Amitriptyline or Nortriptyline (Elavil® or Aventyl HCl®)	\$4 / month – Super-stores drug program

Depression medicines may cause some:

- constipation, diarrhea and nausea
- increased risk of suicidal thoughts and behaviors (18- to 24-year-olds)
- harm to an unborn child
- risk of developing serotonin syndrome, a potentially life-threatening condition
- possible drug-drug interactions

Additional considerations

Category	Medicine (Brand Name)	Additional considerations
SSRIs	Citalopram (Celexa®)	Can cause problems with your heart
	Escitalopram (Lexapro®)	Currently no other issues
	Fluoxetine (Prozac®)	More likely to interact with other drugs you are taking
	Fluvoxamine (Luvox®)	More likely to cause constipation, diarrhea or nausea Not officially recognized as a treatment for Major Depressive Disorder
	Paroxetine (Paxil®)	If you are pregnant, this medicine is more likely to cause problems with your unborn child
	Sertraline (Zoloft®)	More likely to cause diarrhea
SNRIs	Desvenlafaxine (Pristiq®)	Tell your doctor if you have high blood pressure
	Duloxetine (Cymbalta®)	Can help with pain Tell your doctor if you have high blood pressure
	Venlafaxine (Effexor®)	More likely to cause nausea and vomiting Can cause problems with your heart Tell your doctor if you have high blood pressure
Others	Bupropion (Wellbutrin®)	Higher risk of seizures
	Mirtazapine (Remeron®)	Starts to work more quickly
TCAs	Amitriptyline or Nortriptyline (Elavil® or Aventyl HCl®)	More likely to cause constipation, diarrhea or nausea Can help with pain If you are elderly, this medication may not be the best option

A take-home guide to help patients compare depression medicines.

What You Should Know

Will this medicine work for me?

- The antidepressants presented in this decision aid all work the same for treating depression.
- Most people with depression can find one that can make them feel better.
- 6 out of 10 people will feel better with the first antidepressant they try and the rest will have to try other antidepressants before they find the one that is right for them.

How long before I feel better?

- Most people need to take an antidepressant regularly for at least 6 weeks to begin to get the full effect.

Understanding side effects

- Most people taking antidepressants have at least one side effect.
- Many side effects go away after a few weeks, but some only go away after you stop the medicine.

Makers of this aid have no financial relationships with pharmaceutical or device manufacturers.
© 2012 Mayo Foundation for Medical Education and Research. All rights reserved. MC5733-43rev0512

This information reflects the best available research studies. It was prepared by Mayo Clinic researchers without funding from makers of depression medicines.